

COSMOLOGICAL EVOLUTION OF THE UNIVERSE NEUTRAL GAS MASS MEASURED BY QUASAR ABSORPTION SYSTEMS

CELINE PEROUX, RICHARD G. MCMAHON, MIKE IRWIN

Institute of Astronomy, Madingley Road, Cambridge CB3 0HA, UK
e-mail: celine@ast.cam.ac.uk

AND LISA J. STORRIE-LOMBARDI

*SIRTF Science Center, California Institute of Technology, MS 100-22, Pasadena
CA 91125, USA*

The cosmological evolution of neutral hydrogen is an efficient way of tracing structure formation with redshift. It indicates the rate of evolution of gas into stars and hence the gas consumption and rate star formation history of the Universe. In measuring HI, quasar absorbers have proven to be an ideal tool and we use observations from a recent survey for high-redshift quasar absorption systems together with data gathered from the literature to measure the cosmological comoving mass density of neutral gas. This paper assumes $\Omega_M=0.3$, $\Omega_\Lambda=0.7$ and $h=0.65$.

1 Quasar Absorption Systems

Quasar absorbers are systems along the line-of-sight between an observer and a quasar, and thus are observed in absorption in the spectra of the background source. They are classified according to their HI column density: Damped Lyman- α systems (hereafter DLA) have $N(HI) > 10^{20.3}$ atom cm $^{-2}$, Lyman Limit Systems (hereafter LLS) have $N(HI) > 10^{17.2}$ atom cm $^{-2}$ and the Lyman- α forest is composed of systems with smaller column densities. The work presented here is based on observations from a recent high- z survey (Perox et al. 2001) together with a compilation of absorption systems from the literature (Storrie-Lombardi and Wolfe 2000). This included 82 quasars with $z > 4$ and resulted in a sufficiently large sample of quasar absorption systems to enable the study of their properties with redshift (z). Fig 1 shows the cumulative number of quasar absorbers per unit $dX(z)$, for two z ranges, absorbers with $z < 3.5$ and $z > 3.5$, where dX is the distance interval: $X(z) = \int_0^z (1+z)^2 [(1+z)^2(1+z\Omega_M) - z(2+z)\Omega_\Lambda]^{-1/2} dz$. The column density of LLS cannot be derived directly from their equivalent widths as they correspond to the non-linear part of the curve of growth. Nevertheless, the expected number of LLS can be calculated by fitting $N_o(1+z)^\gamma$ to the observed number of systems per unit z and integrating over the z -path surveyed: $LLS_{expected} = \sum_{i=1}^n \int_{z_{min}}^{z_{max}} N_o(1+z)^\gamma dz$. The observations presented




Figure 1. Cumulative number of absorbers per unit distance interval, dX , for two different z ranges. The “steps” correspond to the observed number of DLAs and the data points at $\log N(HI) = 17.2$ are the expected number of LLS derived from the observed number of LLS per unit z . Γ -functions are used for the fits to the data.

in Fig 1 show that there are fewer systems at the highest column density for $z > 3.5$ compared to the lower redshift range, which indicates that we are probing the epoch of assembly of high column density systems from smaller column density units.

2 Cosmological Evolution of Neutral Hydrogen

The neutral gas mass is calculated by integrating the quasar absorbers column density distribution, $f(N, z)$: $\Omega_{HI}(z) = \frac{H_0 \mu m_H}{c \rho_{crit}} \int_{N_{min}}^{\infty} N f(N, z) dN$. We used a Γ -function to fit the data (Fig 1) in order to take into account the HI included below the formal DLA definition. Indeed, DLAs were previously thought to contain the majority of the total HI but our new study shows that 14% at $z < 3.5$ and up to 40% at $z > 3.5$ of the neutral gas is actually contained in smaller column density systems ($10^{17.2} < N(HI) < 10^{20.3}$). The total HI decreases with decreasing z thus matching the pattern expected from gas depletion due to star formation. Details on the method used and more discussion on the results presented here will be available in a subsequent paper.




Figure 2. Cosmological evolution of HI. The circles show the neutral gas contained in DLAs (filled circles are this work, open circles are from Rao and Turnshek 2000 who used HST spectra of quasars with know MgII systems to look for low- z DLAs. Their results are surprisingly high but might be biased by small number statistics or the effect of gravitational lensing.) Vertical error bars correspond to 1σ uncertainties and the horizontal error bars indicate bin sizes. The stars are the total HI including a correction for the neutral gas contained in systems with column densities below the DLA formal definition. The triangle at $z = 0$ is the local HI mass measured by Rao & Briggs 1993. The squares, Ω_{FHP} and Ω_{GO} (Fukugita, Hogan & Peebles 1998 and Gnedin & Ostriker 1992 respectively) are $\Omega_{baryons}$ in local galaxies. Semi-analytical models from Somerville, Primack & Faber 2000 are overplotted.

References

1. Fukugita, M., Hogan, C. and Peebles, P. 1998, ApJ, 503, 518.
2. Gnedin, N. and Ostriker, J., 1992, ApJ, 400, 1.
3. Peroux, C., Storrie-Lombardi, L., McMahon, R., Irwin, M. and Hook, I., 2001, AJ, in press, astro-ph/0101179.
4. Rao, S. and Briggs, F. 1993, ApJ, 419, 515.
5. Rao, S. and Turnshek, D. 2000, ApJS, 130, 1.
6. Somerville, R., Primack, J. and Faber, S. 2000, MNRAS, in press, astro-ph/0006364.
7. Storrie-Lombardi, L. and Wolfe, A. 2000, ApJ, 543, 552.